



**UNIVERSITI PUTRA MALAYSIA**

**SOME ASPECTS OF LARVAL NUTRITION OF THE FRESHWATER  
PRAWN, MACROBRACHIUM ROSENBERGII (DE MAN) WITH AN  
EMPHASIS ON LIPID METABOLISM**

**PAYMOND ROUSTAIA**

**FP 1999 18**

**SOME ASPECTS OF LARVAL NUTRITION OF THE FRESHWATER  
PRAWN, *MACROBRACHIUM ROSENBERGII* (DE MAN) WITH AN  
EMPHASIS ON LIPID METABOLISM**

**BY**

**PAYMON ROUSTAIAN**

**Thesis Submitted in Fulfilment of the Requirements for the  
Degree of Doctor of Philosophy in the Faculty of Agriculture  
Universiti Putra Malaysia**

**December 1999**



To my parents

for their true love, constant trust and principles that guide my life

To my wife

for her devotion, understanding and support during all difficulties

To my children

for making everything worthwhile

and

to many researchers whose works have not given due recognition for the many hours spent in the laboratory and fields to provide humanity with solutions to better life

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in  
fulfilment of the requirements for the degree of Doctor of Philosophy.

**SOME ASPECTS OF LARVAL NUTRITION OF THE FRESHWATER  
PRAWN, *MACROBRACHIUM ROSENBERGII* (DE MAN) WITH AN  
EMPHASIS ON LIPID METABOLISM**

By

**Paymon Roustaian**

**December 1999**

**Chairman : Dr. Mohd Salleh Kamarudin**

**Faculty : Agriculture**

Changes in biochemical composition and fatty acid metabolism during larval development of the freshwater prawn, *Macrobrachium rosenbergii* (de Man) were studied to enhance the current understanding of the physiology and nutritional requirements of the growing larvae. Protein was always the major organic constituent followed by lipids and then carbohydrates. Protein concentration increased during development of both fed and starved larvae, suggesting its important role in morphogenesis. The decline of lipid during the larval development was more rapid for starved larvae which suggests a probable utilisation of lipid as metabolic source of energy. Low levels of carbohydrates in both fed and starved larvae may indicate their limited role in larval metabolism. The amino acid composition appeared to be relatively unchanged during the larval growth, suggesting that the amino acids requirement of the freshwater prawn is relatively constant during larval life. Unlike amino acids, fatty acids

revealed much wider range of variation during larval development. Although a decrease in total saturated fatty acid and monounsaturated fatty acids was observed, polyunsaturated fatty acids tended to increase during larval development. *M. rosenbergii* larvae appeared to have capability to sufficiently convert palmitic (16:0), linoleic (18:2 $n$ -6) and linolenic (18:3 $n$ -3) acids to stearic (18:0), arachidonic (20:4 $n$ -6) and eicosapentaenoic (20:5 $n$ -3) acids, respectively. Findings revealed that dietary lipid level of >15% (dry weight) has detrimental effects on the larval growth and survival. The best performance in terms of postlarval production, dry weight, protein and lipid depositions were observed for the diet containing 12% lipid. The lack of suitability of diets rich in saturated fatty acids for larval freshwater prawn was noted. Despite a wide variation in the dietary saturated to polyunsaturated fatty acid ratio, larvae revealed a narrow range indicative of the importance of such ratio in the larval metabolism. Moreover, regardless of the significant dietary variations in  $n$ -3 to  $n$ -6 ratios, the larval  $n$ -3 to  $n$ -6 ratios were remarkably similar, indicating the importance of such ratio in the larval fatty acid metabolism. The incorporation of linseed oil (as a source of linolenic (18:3 $n$ -3) acid) at 25 to 75 % of supplemented oil increased postlarval production, while larvae fed with linseed oil at 100% revealed significantly lower postlarval production. Although larval tissue level of arachidonic (20:4 $n$ -6) acid was not significantly affected by the dietary level of  $n$ -3 or  $n$ -6 fatty acids, biosynthesis of eicosapentaenoic (20:5 $n$ -3) acid is most likely affected.

Abstrak tesis dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi syarat untuk mendapatkan Ijazah Doktor Falsafah.

**BEBERAPA ASPEK PEMAKANAN LARVA UDANG GALAH,  
*MACROBRACHIUM ROSENBERGII* (DE MAN) DENGAN  
PENEKANAN KE ATAS METABOLISME LIPID**

Oleh

**Paymon Roustaian**

**Disember 1999**

**Pengerusi : Dr. Mohd. Salleh Kamarudin**

**Fakulti : Pertanian**

Perubahan dalam komposisi biokimia dan metabolisme asid lemak semasa perkembangan larva udang galah, *Macrobrachium rosenbergii* (de Man) telah dikaji untuk meningkatkan kefahaman terkini terhadap fisiologi dan keperluan pemakanan larva yang sedang membesar. Protein sentiasa menjadi komposisi organik yang utama diikuti dengan lipid dan karbohidrat. Kepekatan protein meningkat semasa perkembangan kedua-dua larva yang diberi makan dan yang dilaparkan. Ini mencadangkan peranan penting protien dalam morfogenesis. Pengurangan lipid semasa perkembangan larva yang lebih cepat untuk larva yang dilaparkan mencadangkan kemungkinan penggunaan lipid sebagai sumber tenaga. Takat karbohidrat yang rendah di dalam kedua-dua larva yang diberi makan dan yang dilaparkan mungkin menunjukkan peranannya yang terhad dalam metabolisme larva. Komposisi asid amino kelihatan secara relatifnya tidak berubah semasa pertumbuhan larva. Ini mencadangkan bahawa keperluan asid amino udang galah adalah secara relatifnya tetap di sepanjang

peringkat larva. Tidak seperti asid amino, asid lemak menunjukkan variasi yang besar semasa perkembangan larva. Walaupun terdapat pengurangan dalam jumlah asid lemak tepu dan asid lemak mono tak tepu, asid lemak poli tak tepu didapati meningkat semasa perkembangan larva. Larva *M. rosenbergii* kelihatan mempunyai kebolehan untuk menukarkan asid-asid palmetik (16:0), linoleik (18:2 $n$ -6) dan linolenik (18:3 $n$ -3), masing-masingnya, kepada asid-asid stearik (18:0), arakidonik (20:4 $n$ -6) and eikosapentaenoik (20:5 $n$ -3). Keputusan menunjukkan bahawa takat lipid diet melebihi 15% (berat kering) mempunyai kesan yang merencatkan pertumbuhan dan kemandirian larva. Keputusan yang terbaik dari segi pengeluaran pascalarva, berat kering, penghasilan protein dan lipid adalah didapati pada diet yang mengandungi 12% lipid. Secara amnya, didapati bahawa diet yang kaya dengan asid lemak tepu adalah kurang sesuai untuk larva udang galah. Di sebalik variasi yang besar dalam nisbah asid lemak tepu kepada poli tak tepu di dalam diet, tisu larva menunjukkan julat nisbah yang kecil yang mencadangkan kepentingan nisbah tersebut di dalam metabolisme larva. Walaupun terdapat nisbah-nisbah  $n$ -3 :  $n$ -6 dalam diet, nisbah-nisbah  $n$ -3 :  $n$ -6 dalam larva mempunyai persamaan yang menakjubkan. Ini menunjukkan kepentingan nisbah tersebut di dalam metabolisme asid lemak dalam larva. Penambahan minyak "linseed" (sebagai sumber asid linolenik, 18:3 $n$ -3) pada 25 hingga 75% minyak tambahan didapati meningkatkan pengeluaran pascalarva, sementara larva yang diberi 100% minyak "linseed" sebagai minyak tambahan menunjukkan pengurangan yang ketara dalam pengeluaran pascalarva. Walaupun paras asid arakidonik (20:4 $n$ -6) dalam tisu larva tidak terjejas dengan ketaranya dengan takat asid-asid lemak  $n$ -3 atau  $n$ -6 dalam diet, biosintesis asid eikosapentaenoik (20:5 $n$ -3) adalah yang paling berkemungkinan terjejas.

## ACKNOWLEDGEMENT

Words are not enough to express my sincere appreciation for the people who were involved in my training towards obtaining this degree. I am very thankful to my supervisor, Dr Mohd Salleh Kamarudin for his guidance, objective discussions and criticisms, encouragement and support throughout the course of the study. I am also indebted to the members of my supervisory committee, Dr. Hishamuddin Omar, Dr. Che Roos Saad and Dr. Mansor Haji Ahmad for providing some of the facilities and their constructive suggestions.

Thanks are extended to Dr. Mohd Said Saad for providing laboratory facilities during early stages of this study. I am also thankful to technical staff of Hatchery and Pond Complex of UPM, Azmi, Zaidi, Jasni, Nordin and Krishnan for their valuable practical assistance. My appreciation is also extended to Faridah Shamsuddin and Roslan Shariff from UPM Graduate School, and Halim Azizi, Ibrahim Saris and Norlia Jaffar of College Pendita Zaba, and NorHajar AbuBakar and Saheh Said of UPM library for their hospitality and support during my early days in Malaysia. I am also indebted to Dr Usman Inuwasu for teaching me the operation of HPLC.

And finally a special word of thanks to my friends, Hamid Rezai- Marnani and his wife Ellaheh, Annie Christianus and her husband James Lim, Corina and her husband Roland, Ruel Villacorte Eguia, Manuel Laron, Catherlyn Foo, Abeer Alhassan, , Majid Eshagii, Farshad Sheshichian, Abdolsamad Jahangard, Ghada A. Elhag, Wana Amawath, Md. Asaduzzaman, Hamid Khodabakhsh, Gholam Reza Rafiee and Albert Gebrael for their companionship, support and concern.



I certify that an Examination Committee met on 16<sup>th</sup> December 1999 to conduct the final examination of Graduate Student on his Doctor of Philosophy thesis entitled "Some Aspects of Larval Nutrition of the Freshwater Prawn, *Macrobrachium rosenbergii* (de Man) with an emphasis on lipid metabolism" in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. The Committee Members for the candidate are as follows:

Mohd Salleh Kamarudin, Ph.D  
Lecturer  
Faculty of Agriculture  
Universiti Putra Malaysia  
(Chairman)


Hishamuddin Omar, Ph.D  
Lecturer  
Faculty of Science and Environmental Studies  
Universiti Putra Malaysia  
(Member)

Che Roos Saad, Ph.D  
Lecturer  
Institute of Bioscience  
Universiti Putra Malaysia  
(Member)

Mansor Haji Ahmad, Ph.D  
Lecturer  
Faculty of Science and Environmental Studies  
Universiti Putra Malaysia  
(Member)

Mohd Ridzwan Abd. Halim, Ph.D  
Associate Professor  
Faculty of Agriculture  
Universiti Putra Malaysia  
(Independent Examiner)

Santosh P. Lall, Ph.D  
Professor  
Dalhousie University  
(Independent Examiner)



MOHD. GHAZALI MOHAYIDIN, PH.D,  
Professor/Deputy Dean of Graduate School,  
Universiti Putra Malaysia

Date: 18 JAN 2000

This Thesis submitted to the Senate of Universiti Putra Malaysia and was accepted as fulfilment of the requirements for the degree of Doctor of Philosophy.



---

KAMIS AWANG, Ph.D,  
Associate Professor  
Dean of Graduate school,  
Universiti Putra Malaysia

Date: **10 FEB 2000**

---

## DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.



---

Paymon Roustaian  
December 1999

## TABLE OF CONTENTS

	Page
<b>DEDICATION.....</b>	ii
<b>ABSTRACT.....</b>	iii
<b>ABSTRAK.....</b>	v
<b>ACKNOWLEDGMENTS.....</b>	vii
<b>APPROVAL SHEETS</b>	viii
<b>DECLARATION FORM</b>	x
<b>LIST OF TABLES.....</b>	xiv
<b>LIST OF FIGURES.....</b>	xvii
<b>LIST OF ABBREVIATIONS.....</b>	xviii
<b>CHAPTER</b>	
<b>I INTRODUCTION.....</b>	1
Background of the study.....	1
Statement of the problem.....	4
Significance of the study.....	6
Objectives	8
<b>II LITERATURE REVIEW.....</b>	10
Egg hatching and larval production of <i>M. rosenbergii</i> .....	11
Hatchery systems for rearing larval <i>M. rosenbergii</i> .....	13
<i>M. rosenbergii</i> larval rearing operation.....	15
Feed and feeding behavior of larval <i>M. rosenbergii</i> .....	18
Nutritional requirements of <i>M. rosenbergii</i> larvae.....	20
Protein and amino acid requirements.....	20
Lipid and fatty acid requirements.....	21
Carbohydrates requirements.....	24
Energy requirements and sources.....	25
Replacement of <i>Artemia</i> in the larval culture of <i>M. rosenbergii</i> .....	26
<b>III GENERAL METHODOLOGY.....</b>	28
Larval management.....	28
Larval rearing.....	28
<i>Artemia</i> preparation.....	30
Larval Staging Index.....	30
Water quality monitoring.....	30
Biochemical evaluation of the larvae.....	31
Sampling.....	31
Determination of the soluble protein.....	31
Determination of the soluble carbohydrates.....	31
Determination of the total lipids.....	32
Amino acid analysis.....	32
Fatty acid analysis.....	33



<b>IV</b>	<b>CHANGES IN PROTEIN, LIPID, AND CARBOHYDRATE CONCENTRATION DURING LARVAL DEVELOPMENT OF <i>MACROBRACHIUM ROSENBERGII</i>.....</b>	<b>35</b>
	Materials and Methods.....	36
	Results.....	38
	Discussion.....	47
<b>V</b>	<b>CHANGES IN AMINO ACIDS AND FATTY ACIDS PROFILE DURING LARVAL DEVELOPMENT OF FRESHWATER PRAWN, <i>MACROBRACHIUM ROSENBERGII</i> (DE MAN).....</b>	<b>53</b>
	Materials and Methods.....	53
	Results.....	54
	Discussion.....	66
	Amino acids.....	66
	Fatty acids.....	68
<b>VI</b>	<b>WEANING OF <i>MACROBRACHIUM ROSENBERGII</i> LARVAE FROM <i>ARTEMIA</i> TO AN ARTIFICIAL DIET.....</b>	<b>74</b>
	Materials and Methods.....	75
	Results.....	77
	Discussion.....	81
<b>VII</b>	<b>THE EFFECTS OF DIETARY LIPID LEVEL ON THE <i>MACROBRACHIUM ROSENBERGII</i> LARVAL PERFORMANCE, POSTLARVAL PRODUCTION AND BIOCHEMICAL COMPOSITION.....</b>	<b>85</b>
	Materials and Methods.....	86
	Results.....	88
	Discussion.....	92
<b>VIII</b>	<b>THE EFFECTS OF DIETARY LIPID SOURCES ON THE <i>MACROBRACHIUM ROSENBERGII</i> LARVAL PERFORMANCE, POSTLARVAL PRODUCTION AND FATTY ACID COMPOSITION.....</b>	<b>95</b>
	Materials and Methods.....	95
	Results.....	98
	Discussion.....	104

<b>IX</b>	<b>THE EFFECTS OF DIFFERENT DIETARY RATIOS OF CORN OIL/LINSEED OIL MIXTURE ON THE <i>MACROBRACHIUM ROSENBERGII</i> LARVAL PERFORMANCE, POSTLARVAL PRODUCTION AND FATTY ACID COMPOSITION.....</b>	<b>107</b>
	Materials and Methods.....	108
	Results.....	113
	Discussion.....	116
<b>X</b>	<b>GENERAL DISCUSSION, CONCLUSIONS AND RECOMMENDATIONS.....</b>	<b>121</b>
	General Discussion.....	121
	Conclusions.....	128
	Recommendations.....	129
	<b>BIBLIOGRAPHY.....</b>	<b>130</b>
	<b>APPENDIX.....</b>	<b>143</b>
	<b>VITA.....</b>	<b>145</b>

## LIST OF TABLES

Table	Page
1.1 Examples of value of farmed freshwater prawns adapted from New (1995).....	3
2.1 Optimal characteristics of important physicochemical features in larviculture of <i>M. rosenbergii</i> .....	17
2.2 Fatty acid profiles of some freshwater and marine crustaceans .....	23
2.3 Trials on replacement of <i>Artemia</i> in <i>M. rosenbergii</i> larviculture.....	27
4.1 Carbohydrate moieties (% dry weight) during larval development of <i>M. rosenbergii</i> reared under fed and starved conditions.....	46
4.2 Caloric content of <i>M. rosenbergii</i> during larval development as calculated from specific enthalpies of combustion.....	48
5.1 Essential amino acid profile (% of total amino acids) of <i>M. rosenbergii</i> at different stages of development.....	55
5.2 Non essential amino acid profile (% of total amino acids) of <i>M. rosenbergii</i> at different stages of development.....	56
5.3 Essential amino acid ratio (A/E) of <i>M. rosenbergii</i> at different stages of development.....	57
5.4 Fatty acid composition (percent of total FAME) of larval stages of <i>M. rosenbergii</i> and newly hatched <i>Artemia</i> nauplii (Aquafauna BioMarine Inc., Calif).....	59
5.5 Total lipid (ng per µg dry wt) and fatty acid composition (ng fatty acid per µg dry wt.) of larval stages of <i>M. rosenbergii</i> and newly hatched <i>Artemia</i> nauplii (Aquafauna BioMarine Inc., Calif).....	60
5.7 Comparison of fatty acid composition (percent of total FAME) of broodstock eggs (orange) under different environments (Das, 1995) and first larval stage of <i>M. rosenbergii</i> .....	69

6.1	Different feeding schemes employed in gradual weaning of <i>M. rosenbergii</i> larvae from <i>Artemia</i> to an artificial diet.....	76
6.2	General composition and chemical analysis of test diets..	78
6.3	Production of <i>M. rosenbergii</i> postlarvae from various weaning schemes for a period of 42 days.....	79
7.1	General composition and chemical analysis of test diets..	87
7.2	Effects of dietary lipid levels on survival, postlarvae production, dry weight, protein and lipid deposition of the surviving larvae/postlarvae <i>M. rosenbergii</i> .....	91
8.1	General composition and chemical analysis of test diets..	97
8.2	Fatty acid composition (percent of total FAME) of the different dietary lipid sources used in the experiment.....	99
8.3	Fatty acid composition (percent of total FAME) of the different diets and <i>Artemia</i> nauplii (Microfeast, Oklahoma, USA) used in the experiment.....	100
8.4	Growth response and biochemical composition of the the surviving larvae/postlarvae <i>M. rosenbergii</i> fed with diets containing different sources of lipids.....	101
8.5	Fatty acid composition (percent of total FAME) of the surviving larvae/postlarvae <i>M. rosenbergii</i> fed with diets containing different lipid sources.....	103
9.1	General composition and chemical analysis of test diets...	110
9.2	Fatty acid composition (percent of total FAME) of different mixture of the corn oil/ linseed oil used in the trial. Data are the mean values of three determinations...	111
9.3	Fatty acid composition (percent of total FAME) of different dietary treatments used in the trial.....	112
9.4	Effects of compounded diets containing different ratio of corn oil / linseed oil on postlarval (PL) production, dry weight, protein and lipid deposition of the surviving larvae/postlarvae <i>M. rosenbergii</i> .....	114



9.5	Fatty acid composition (percent of total FAME) of the surviving larvae/postlarvae <i>M. rosenbergii</i> fed with diets containing different ratio of corn oil/linseed mixture.....	115
-----	--	-----

## LIST OF FIGURES

Figures	Page
4.1 Survival of <i>M. rosenbergii</i> larvae reared under fed and starved conditions.....	39
4.2 Larval growth (expressed as MDS) of <i>M. rosenbergii</i> reared under fed and starved conditions.....	40
4.3 Fresh weight ( $\mu\text{g}$ /larva) of <i>M. rosenbergii</i> larvae at different stages of development reared under fed and starved conditions.....	41
4.4 Dry weight ( $\mu\text{g}$ /larva) of <i>M. rosenbergii</i> larvae at different stages of development reared under fed and starved conditions.....	42
4.5 Biochemical composition of <i>M. rosenbergii</i> larvae at different stages of development reared under fed (with <i>Artemia</i> ) condition.....	44
4.6 Biochemical composition of <i>M. rosenbergii</i> larvae at different stages of development reared under starved condition.....	45
5.1 Percent change of major fatty acid classes related to previous larval stage expressed per unit dry weight during larval development of <i>M. rosenbergii</i> .....	64
5.2 Percent change of major fatty acid classes related to previous larval stage expressed per larva during larval development of <i>M. rosenbergii</i> .....	65
6.1 Survival of <i>M. rosenbergii</i> larvae reared under various weaning schemes.....	80
6.2 Larval growth (expressed as MDS) of <i>M. rosenbergii</i> reared under various weaning schemes.....	82
7.1 Survival of <i>M. rosenbergii</i> larvae fed with diets containing different lipid levels.....	89
7.2 Larval growth (expressed as MDS) of <i>M. rosenbergii</i> fed with diets containing different lipid levels.....	90
10.1 Changes in <i>n</i> -3 to <i>n</i> -6 ratio during various life stages of <i>M. rosenbergii</i> . LL=Late Larvae, EPL=Early postlarvae..	125



## LIST OF ABBREVIATIONS

A/E = Essential amino acid ratio

ANOVA = Analysis of variance

CGAF = Calories per gram ash-free dry weight

CLO = Cod liver oil

DW = Dry weight

EPL= Early postlarvae

FAME = Fatty acid methyl esters

HUFA = Highly unsaturated fatty acids

LL= Late larvae

LSD = Least significant difference

MDS = Mean developmental stage

MUFA = Monounsaturated fatty acids

PL = Postlarval

ppm = Part per million

ppt = Part per thousand

PUFA = Polyunsaturated fatty acids

sd = Standard deviation

SFA = Saturated fatty acids

# **CHAPTER I**

## **INTRODUCTION**

### **Background of the Study**

As capture fishery approaches its maximum sustainable yield, the significance of aquaculture to provide nutritious food of high protein value is gaining more appreciation at global spectrum. Moreover, economic restriction such as increasing operational cost of fishing fleets as well as diminishing and unpredictable yields of oceans will most probably contribute to aquaculture growth and expansion in years to come.

Information compiled by tremendous research efforts and its application at various aspects of aquatic farming both in freshwater and marine environment have profoundly altered farming of aquatic organisms during the latter part of 20<sup>th</sup> century. The extent of this advancement has been as such that nowadays it is referred to as the “blue revolution” (Lovell, 1989). Aquaculture is now recognized as a viable and a profitable practice that has attracted commercial investment at a large scale.

Farming of aquatic crustaceans has become increasingly significant due to their superior taste, high nutritious values and excellent market (New, 1995). The global production of cultured crustaceans in late 80's/early 1990's was approximately 982,000 metric tones annually, which contributed to about 20-25% of world crustacean supplies (Lee and Wickins, 1992; New, 1995). Out of this figure, shrimps

as referred to *Penaeus* and *Metapenaeus* spp. according to FAO terminology (Fast, 1992) provided 90% while prawns (*Macrobrachium* spp. according to FAO terminology) contributed 3.2%. Accordingly, crayfish and other crustaceans contributed to 6.8% of total farmed crustacean production (New, 1995).

Despite the apparent dominance of penaeid farming at global scope, an increasing trend in freshwater prawn production and its recent exports from some Southeast Asian countries have demonstrated an ample culture potential of this commodity. Farmed prawn production globally expanded by a factor of 3.5 in the seven years period from 1984 (~10500 metric tonnes) to 1991 (~ 37000 metric tonnes) (New, 1995). Prawn production in Malaysia also revealed an increasing trend from 68 metric tonnes in 1984, and 78.5 tonnes in 1995 to 95.6 tones in 1996 (Anon, 1985, 1996, 1998). Nearly all records of farmed prawn are attributed to *Macrobrachium rosenbergii* which is one of the largest species of the *Macrobrachium* genus (New, 1995). Although the genus possesses some other large species of economic value, culture of *M. rosenbergii* is favored due to its good survival and rapid growth (New, 1990), resistance to common disease problems (Johnson, 1982), wide range of salinity and temperature tolerance of postlarvae and juveniles (Smith *et al.*, 1982), ability to become mature, mate and spawn repeatedly in captivity (Wickins, 1976) and omnivorous feeding regime (Lee *et al.*, 1980; Weidenback, 1982). *M. rosenbergii* is also suitable to culture in inland freshwater bodies which may reduce pressure on limited coastal resources and increase income generating activities. These factors plus high market price at global scale (Table 1.1) render the *M. rosenbergii* culture a viable practice (New, 1995). However, large-scale production

Table 1.1: Examples of value of farmed freshwater prawns adapted from New (1995).

Location	Date	Price	
		(US\$ / kg)	Note
Brazil	1991	9-12	Whole; on ice, semi-wholesale to restaurants
	1993	5-12	Whole; domestic
France	1987	10	Imported from south-east Asia
		16	Imported from French Caribbean
	1989	20	Imported from French Caribbean
Martinique	1993	23-30	(20-30/kg) Whole; fresh or live
		19-23	(30-40/kg) Whole; fresh or live
Nepal	1994	6-7	Whole
Taiwan	1993	12.6	Domestic; Whole
United Kingdom	1994	18.2	Heads-off, shell on (supermarket)
USA	1994	22-27	Live (farm-gate)
Malaysia	1999	7-8	Whole; domestic (personal observation)

of *M. rosenbergii* has yet to be refined in terms of seed production and appropriate culture technology for rural farmers.

### **Statement of the problem**

The high market value and the great demand of *M. rosenbergii* in Malaysia provide a strong stimulus for its culture. Although the larval rearing of *M. rosenbergii* was first reported in Malaysia, its culture industry has not developed as expected. Shortage of seeds, absence of a suitable feed, absence of an appropriate culture system and poor management are recognized as the major bottlenecks in development of culture industry in this country (Ang *et al.*, 1990).

To insure a stable local or international market, a reliable, constant and reasonable quantities of high quality cultured prawn is a prerequisite. This, in turn, is ultimately linked to status of juvenile production which is the corner stone in culture of any aquatic species. Freshwater prawn juveniles can either be obtained from wild or through hatcheries. Collection of juveniles from natural sources for stocking in ponds is faced with many problems such as difficulties in obtaining individuals of the same uniform sizes, limited and unreliable supply, laborious identification, difficulties and high cost associated with transport of juveniles from collection sites to growing farms (Bian and Pang, 1982; Suharto *et al.*, 1982). A well-operated prawn hatchery is thus the best alternative as the source of seed supply.

Hatchery larviculture and production of postlarvae of *M. rosenbergii*, in turn, are far from perfect and require much improvement. In a typical *M. rosenbergii* larviculture, mortality is caused either by poor water quality, problems at first

feeding, disease or any combination of such factors (New, 1982). These setbacks, in general, stem out from a prolonged larval life of *M. rosenbergii* (Wickin, 1976; New, 1990).

Among the problems associated with a long larviculture period as well as the expenses concerning physical maintenance and labor, the dependence of larvae upon *Artemia* nauplii as indispensable food (Murai and Andrews, 1978; Aniello and Singh, 1982; Lovett and Felder, 1988; Alam *et al.*, 1993a) is considered a major parameter that restricts the development of this industry in developing nations. Despite the ease of use and preparation as well as its nutritional qualities, the main disadvantage of *Artemia* based diet in larviculture of freshwater prawn is its high cost and occasional scarcity especially in developing countries. Cost of feeding may well reach to as high as 60% of the total production costs in *M. rosenbergii* hatcheries (Hagood and Willis, 1976). Variations in hatchability and nutritional value of *Artemia* from different geographic sources and even from one batch to another within the same strain, are also of great concern in the hatcheries operation (Daniels *et al.*, 1992; Sorgeloos and Leger, 1992). Another disadvantage of *Artemia* based diet is that brine shrimp exuvia and shed cyst capsules accumulates in larval culture vessels. Bacterial degradation of these materials fouls the water while accumulated debris entangles larvae and increases larval mortality (Lovett and Felder, 1988). Of parallel importance is the nutritional value of *Artemia* nauplii that apparently decreases rapidly after hatching due to reduced yolk material (Maddox and Manzi, 1976).

Substitution by any relevantly non-expensive live food could to some extent alleviate the problem. However, the number of undesirable side effects, such as man



power, required skill, nutritional variability and high risk of pathogenic infection associated with use of live food in general, are limiting parameters in application of such feed. The development and application of high quality artificial diet that can reduce the technical difficulties and operational costs associated with live feed preparation would be highly advantageous.

Problems associated with digestibility, acceptability, stability and nutritional status of the formulated diets may single-handedly or in combination contribute to lack of success in total replacement with artificial diet in caridean shrimps larval culture including *M. rosenbergii* (Jones *et al.* 1993; New, 1995). The most promising results for *M. rosenbergii* larvae were obtained by Deru (1990) who reared the larvae using Frippak microencapsulated diets exclusively from day 16 (larval stage of 6-7) to metamorphosis. However, high cost of this diet in developing countries would still prevent the growth of freshwater prawn aquaculture industry. An easily available, cheap, and nutritionally sound artificial diet is therefore desired to reduce the dependency of *Artemia* in the larval rearing of *M. rosenbergii* in developing nations. Understanding of biochemical and physiological processes pertaining to larval *M. rosenbergii* life cycle would provide valuable information in formulating such diet.

### Significance of the study

As it was indicated earlier, the major challenge in the development of culture system for larval *M. rosenbergii* is to reduce larval dependency upon supplies of live food organisms which are both costly and nutritionally inconsistent. The development of high-quality artificial diet could reduce the high cost of live food as well as ameliorate water quality and disease problems (Bengtson, 1993). Knowledge